



HP Xenon by d-DNP using the clinical GE SPINlab polarizer system

Mariager, C. ; Ringgaard, S. ; Ardenkjær-Larsen, Jan Henrik; Laustsen, C.

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Mariager, C., Ringgaard, S., Ardenkjær-Larsen, J. H., & Laustsen, C. (2016). *HP Xenon by d-DNP using the clinical GE SPINlab polarizer system*. Abstract from EUROMAR 2016, Aarhus, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

HP Xenon by d-DNP using the clinical GE SPINlab polarizer system

C. Mariager,^{*} S. Ringgaard,[‡] J.H. Ardenkjaer-Larsen^{**,†} and C. Laustsen[‡]

^{*}Department of Physics and Astronomy, Aarhus University, [‡]Department of Clinical Medicine, Aarhus University Hospital, ^{**}Department of Electrical Engineering, Technical University of Denmark, [†]GE Healthcare, Denmark

Purpose: Hyperpolarized (HP) ^{129}Xe for magnetic resonance (MR) has wide use in material sciences, structural chemistry and medicine. In the medical field ^{129}Xe has been demonstrated as a useful probe for MR lung imaging and proposed as a blood tracer for *in vivo* perfusion imaging¹. The Xenon probe is therefore promising for clinical assessment of brain function², and may aid in the diagnosis and prognosis of brain diseases. Standard production of HP ^{129}Xe is via spin exchange optical pumping (SEOP). However, reports of large polarization enhancements for ^{129}Xe via dynamic nuclear polarisation (DNP)^{1,3} have raised expectations that DNP can be an alternative method for producing HP ^{129}Xe for patients. We therefore investigated the possibility for production of HP ^{129}Xe using the clinical GE SPINlab polarizer, extending the practical use of the system.

Materials and methods: Solid state samples of 0.80 ± 0.01 mL 1-Propanol / 12 mM AH111501 radical, infused with natural abundance (n.a.) or enriched (70% ^{129}Xe) Xenon gas at 5.0 ± 0.2 bar partial pressure for 60 ± 1 minutes at room temperature, were prepared and inserted into the 5 T SPINlab polarizer. Microwaves at 140.01 GHz were used to irradiate the samples for 150 ± 2 minutes at 0.90 ± 0.04 K followed by dissolution, transport and MR measurement. HP / thermal MR experiments were performed ($n = 3 / 2$) with $9^\circ \pm 1^\circ / 90^\circ \pm 1^\circ$ flip angles and $\text{TR} = 1 \text{ s} / 300$

s on a 9.4 T small bore rodent MR scanner using a volume radio frequency (RF) coil with 1 / 200 averages. Transport times for the n.a. / enriched gas experiments were $32 \pm 1 \text{ s} / 27 \pm 1 \text{ s}$ at earth field.

Results: The experiments resulted in enhancements / polarizations of $214 \pm 22 / 0.18 \pm 0.02 \%$ and $431 \pm 30 / 0.37 \pm 0.03 \%$ with $T_1 = 29 \pm 2 \text{ s}$ and $26 \pm 2 \text{ s}$ for n.a. and enriched gas, respectively.

Conclusion: The experiments have shown that it is possible to hyperpolarize ^{129}Xe using the SPINlab system. However, substantial work is required to achieve similar enhancements to SEOP. Strides should also be taken to prolong the T_1 lifetime of the resulting HP gas. These issues can be addressed on multiple fronts, including optimization of the radical and Xenon concentrations, using different solvents and transporting the HP gas in a dedicated electromagnetic carrier.




Figure 1: Decay of one of the recorded enriched HP ^{129}Xe spectra after phasing and exponential line broadening.

¹ A. Capozzi, C. Roussel, A. Comment, J.N. Hyacinthe; Optimal Glass-Forming Solvent Brings Sublimation Dynamic Nuclear Polarization to ^{129}Xe Hyperpolarization Biomedical Imaging Standards. *Phys. Chem. C* 199 (2015) 5020-5025.

² M. L. Mazzanti, R. P. Walvick, X. Zhou, Y. Sun, N. Shah, J. Mansour, J. Gereige, M. S. Albert; Distribution of Hyperpolarized Xenon in the Brain Following Sensory Stimulation: Preliminary MRI Findings. *Plos One*. 7 (2011) 1-7.

³ A. Comment, S. Jannin, J. Hyacinthe, P. Miville, R. Sarkar, P. Ahuja, P. R. Vasos, X. Montet, F. Lazeyras, J. Valle, P. Hautle, J. A. Konter, B. van den Brandt, J. Ansermet, R. Gruetter, and G. Bodenhausen; Hyperpolarizing Gases via Dynamic Nuclear Polarization and Sublimation. *Phys. Rev. Lett.* 105 (2010) 018104.